

ZRSR2 Antibody

PACO48590

Description

This ZRSR2 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO48590
Contents:	50µg Bradford Reagent: 1 vial (2ml)
Category:	Polyclonal Antibody
Synonyms:	CCCH type zinc finger antibody, CCCH type zinc finger, RNA binding motif and serine/arginine rich protein 2 antibody, MGC142014 antibody, MGC142040 antibody, Renal carcinoma antigen NY REN 20 antibody, Renal carcinoma antigen NY-REN-20 antibody, RNA-binding motif and serine/arginine rich protein 2 antibody, U2 small nuclear ribonucleoprotein auxiliary factor 35 kDa subunit related protein 2 antibody, U2 small nuclear ribonucleoprotein auxiliary factor 35 kDa subunit-related protein 2 antibody, U2(RNU2) small nuclear RNA auxiliary factor 1 like 2 antibody, U2(RNU2) small nuclear RNA auxiliary f... Read more
Clone:	Polyclonal
Applications:	ELISA IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q15696
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human U2 small nuclear ribonucleoprotein auxiliary factor 35 kDa subunit-related protein 2 protein (191-372AA)
Immunogen Species:	Homo sapiens (Human)

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS,
PH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:20-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

Immunohistochemistry of paraffin-embedded human testis tissue using PACO48590 at dilution of 1:100