

KDM6A Antibody

PACO55826

Description

This KDM6A Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO55826
Contents:	50µg Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	bA386N14.2 (ubiquitously transcribed X chromosome tetratricopeptide repeat protein (UTX) antibody, bA386N14.2 antibody, bA386N14.2 ubiquitously transcribed X chromosome tetratricopeptide repeat protein UTX antibody, DKFZp686A03225 antibody, Histone demethylase UTX antibody, KABUK2 antibody, Kdm6a antibody, KDM6A_HUMAN antibody, Lysine (K) specific demethylase 6A antibody, Lysine K specific demethylase 6A antibody, Lysine specific demethylase 6A ubiquitously transcribed tetratricopeptide repeat gene X chromosome antibody, Lysine-specific demethylase 6A antibody, MGC141941 antibody, Ubiquitously...Read more
Clone:	Polyclonal
Applications:	ELISA IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	O15550
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human Lysine-specific demethylase 6A protein (620-749AA)
Immunogen Species:	Homo sapiens (Human)

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:200-1:500

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

IHC image of PACO55826 diluted at 1:400 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.