

## LAPTM4A Antibody

PACO58164

### Description

---

This LAPTM4A Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

|                      |   |
|----------------------|---|
| <b>SKU:</b>          | PACO58164   |
| <b>Contents:</b>     | 50µg<br>Bradford Reagent: 1 vial (2ml)  |
| <b>Category:</b>     | -   |
| <b>Synonyms:</b>     | LAPTM4A, KIAA0108, LAPTM4, MBNT, MTRP, UNQ1846/PRO3574, Lysosomal-associated transmembrane protein 4A, Golgi 4-transmembrane-spanning transporter MTP |
| <b>Clone:</b>        | Polyclonal  |
| <b>Applications:</b> | <a href="#">ELISA</a> <a href="#">IHC</a> <a href="#">IF</a>  |
| <b>Conjugation:</b>  | Non-conjugated  |
| <b>Reactivity:</b>   | Human   |

### Antibody Data

---

|                           |   |
|---------------------------|---|
| <b>Isotype:</b>           | IgG   |
| <b>Uniprot:</b>           | Q15012  |
| <b>Host Species:</b>      | Rabbit  |
| <b>Purification:</b>      | >95%, Protein G purified  |
| <b>Immunogen:</b>         | Recombinant Human Lysosomal-associated transmembrane protein 4A protein (181-233AA) |
| <b>Immunogen Species:</b> | Homo sapiens (Human)  |
| <b>Buffer:</b>            | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4       |
| <b>Form:</b>              | Liquid  |

## Preparation & Storage


**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

| Recommended Dilutions: | Application | Recommended Dilution |
|------------------------|-------------|----------------------|
|                        | IHC         | 1:200-1:500          |
|                        | IF          | 1:50-1:200           |

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


## Validation Data

### Image




### Description

IHC image of PACO58164 diluted at 1:300 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of PACO58164 diluted at 1:300 and staining in paraffin-embedded human cervical cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HepG2 cells with PACO58164 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).