

BMPER Antibody

PACO58344

Description

This BMPER Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO58344
Contents:	50µg Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	BMP binding endothelial regulator antibody, BMP binding endothelial regulator precursor protein antibody, BMP-binding endothelial regulator protein antibody, BMPER antibody, BMPER_HUMAN antibody, Bone morphogenetic protein binding endothelial cell precursor derived regulator antibody, Bone morphogenetic protein-binding endothelial cell precursor-derived regulator antibody, CRIM3 antibody, Crossveinless 2 antibody, CV 2 antibody, CV2 antibody, hCV2 antibody, Protein crossveinless 2 antibody, Protein crossveinless-2 antibody
Clone:	Polyclonal
Applications:	ELISA IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q8N8U9
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human BMP-binding endothelial regulator protein (271-418AA)
Immunogen Species:	Homo sapiens (Human)

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:200-1:500

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

IHC image of PACO58344 diluted at 1:200 and staining in paraffin-embedded human pancreatic cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.