

UMODL1 Antibody

PACO59932

Description

This UMODL1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO59932
Contents:	50µg Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	D17Ert488e antibody, Olfactorin antibody, OTTHUMP00000109312 antibody, OTTHUMP00000109313 antibody, RGD1566039 antibody, UMODL1 antibody, UROL1_HUMAN antibody, Uromodulin-like 1 antibody
Clone:	Polyclonal
Applications:	ELISA WB IHC IF
Conjugation:	Non-conjugated
Reactivity:	Human, Mouse

Antibody Data

Isotype:	IgG
Uniprot:	Q5DID0
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human Uromodulin-like 1 protein (537-654AA)
Immunogen Species:	Homo sapiens (Human)
Buffer:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Form:	Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	WB	1:500-1:5000
	IHC	1:200-1:500
	IF	1:50-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data

Image




Description

Western Blot Positive WB detected in: NIH/3T3 whole cell lysate, MCF-7 whole cell lysate All lanes: UMODL1 antibody at 5.1 μ g/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 145, 157, 137, 150 kDa Observed band size: 145, 157 kDa




IHC image of PACO59932 diluted at 1:300 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of PACO59932 diluted at 1:300 and staining in paraffin-embedded human prostate cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary

antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with PACO59932 at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).