

PSG9 Antibody

PACO62023

Description

This PSG9 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO62023
Contents:	50µl Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	pregnancy specific beta 1 glycoprotein (PSG) antibody, pregnancy specific beta 1 glycoprotein 11 antibody, pregnancy specific beta 1 glycoprotein 9 antibody, pregnancy specific beta 1 glycoprotein B antibody, Pregnancy-specific beta-1 glycoprotein B antibody, Pregnancy-specific beta-1-glycoprotein 11 antibody, Pregnancy-specific beta-1-glycoprotein 9 antibody, Pregnancy-specific glycoprotein 11 antibody, Pregnancy-specific glycoprotein 7 antibody, Pregnancy-specific glycoprotein 9 antibody, PS beta B antibody, PS-beta-B antibody, PS-beta-G-11 antibody, PS-beta-G-9 antibody, PS34 antibody, PSBG...Read more
Clone:	Polyclonal
Applications:	ELISA IHC IF
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q00887
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human Pregnancy-specific beta-1-glycoprotein 9 protein (241-426AA)
Immunogen Species:	Homo sapiens (Human)

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:20-1:200
	IF	1:50-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data


Image



Description

IHC image of PACO62023 diluted at 1:100 and staining in paraffin-embedded human liver cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

IHC image of PACO62023 diluted at 1:100 and staining in paraffin-embedded human placenta tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with PACO62023 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).