

USP31 Antibody

PACO62495

Description

This USP31 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO62495
Contents:	50µl Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	Deubiquitinating enzyme 31 antibody, KIAA1203 antibody, Ubiquitin carboxyl terminal hydrolase 31 antibody, Ubiquitin carboxyl-terminal hydrolase 31 antibody, Ubiquitin specific peptidase 31 antibody, Ubiquitin specific processing protease 31 antibody, Ubiquitin specific protease 31 antibody, Ubiquitin specific proteinase 31 antibody, Ubiquitin thioesterase 31 antibody, Ubiquitin thiolesterase 31 antibody, Ubiquitin-specific-processing protease 31 antibody, UBP31_HUMAN antibody, USP 31 antibody, USP31 antibody
Clone:	Polyclonal
Applications:	ELISA IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q70CQ4
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Recombinant Human Ubiquitin carboxyl-terminal hydrolase 31 protein (1069-1196AA)
Immunogen Species:	Homo sapiens (Human)

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:200-1:500

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

IHC image of PACO62495 diluted at 1:300 and staining in paraffin-embedded human adrenal gland tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.