

WFIKKN1 Antibody

PACO63183

Description

This WFIKKN1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO63183
Contents:	50µl Bradford Reagent: 1 vial (2ml)
Category:	-
Synonyms:	WFIKKN1 antibody, C16orf12 antibody, GASP2 antibody, WFIKKN antibody, PSEC0040 antibody, WAP antibody, Kazal antibody, immunoglobulin antibody, Kunitz and NTR domain-containing protein 1 antibody, Growth and differentiation factor-associated serum protein 2 antibody, GASP-2 antibody, hGASP-2 antibody, WAP antibody, follistatin antibody, immunoglobulin antibody, Kunitz and NTR domain-containing protein antibody
Clone:	Polyclonal
Applications:	ELISA IHC
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q96NZ8
Host Species:	Rabbit
Purification:	>95%, Protein G purified
Immunogen:	Peptide sequence from Human WAP, Kazal, immunoglobulin, Kunitz and NTR domain-containing protein 1 protein (187-207AA)
Immunogen Species:	Homo sapiens (Human)

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IHC	1:200-1:500

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

IHC image of PACO63183 diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.