

WDR73 Antibody

PACO63427

Description

This WDR73 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

| | |
|----------------------|---|
| SKU: | PACO63427 |
| Contents: | 50µl Bradford Reagent: 1 vial (2ml) |
| Category: | - |
| Synonyms: | WDR73 antibody, HSPC264WD repeat-containing protein 73 antibody |
| Clone: | Polyclonal |
| Applications: | ELISA WB IHC IF |
| Conjugation: | Non-conjugated |
| Reactivity: | Human |

Antibody Data

| | |
|---------------------------|---|
| Isotype: | IgG |
| Uniprot: | Q6P4I2 |
| Host Species: | Rabbit |
| Purification: | Antigen Affinity Purified |
| Immunogen: | Recombinant Human WD repeat-containing protein 73 protein (163-366AA) |
| Immunogen Species: | Homo sapiens (Human) |
| Buffer: | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Form: | Liquid |

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Store Bradford Reagent at Room Temperature for 1 Year.

| Recommended Dilutions: | Application | Recommended Dilution |
|------------------------|-------------|----------------------|
| | WB | 1:500-1:2000 |
| | IHC | 1:20-1:200 |
| | IF | 1:50-1:200 |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol


Validation Data

Image




Description

Western Blot Positive WB detected in: MCF-7 whole cell lysate, U87 whole cell lysate, 293 whole cell lysate, A549 whole cell lysate, THP-1 whole cell lysate All lanes: WDR73 antibody at 1:1000 Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 42 kDa Observed band size: 42 kDa



IHC image of PACO63427 diluted at 1:100 and staining in paraffin-embedded human endometrial cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of SH-SY5Y cells with PACO63427 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).