

SLC39A10 Antibody

PACO63519

Description

This SLC39A10 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	PACO63519
Contents:	50µl Bradford Reagent: 1 vial (2ml)
Category:	Polyclonal Antibody
Synonyms:	2900042E17Rik antibody, 5430433I10 antibody, DKFZp781L10106 antibody, KIAA1265 antibody, LZT-Hs2 antibody, MGC126565 antibody, MGC138428 antibody, mKIAA1265 antibody, OTTMUSP00000023810 antibody, S39AA_HUMAN antibody, slc39a10 antibody, Solute carrier family 39 (metal ion transporter), member 10 antibody, Solute carrier family 39 (zinc transporter), member 10 antibody, Solute carrier family 39 member 10 antibody, Zinc transporter ZIP10 antibody, ZIP-10 antibody, ZIP10 antibody, Zrt- and Irt-like protein 10 antibody
Clone:	Polyclonal
Applications:	ELISA IF
Conjugation:	Non-conjugated
Reactivity:	Human

Antibody Data

Isotype:	IgG
Uniprot:	Q9ULF5
Host Species:	Rabbit
Purification:	Antigen Affinity Purified
Immunogen:	Recombinant Human Zinc transporter ZIP10 protein (41-317AA)
Immunogen Species:	Homo sapiens (Human)

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Buffer: Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Form: Liquid

Preparation & Storage


Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Store Bradford Reagent at Room Temperature for 1 Year.

Recommended Dilutions:	Application	Recommended Dilution
	IF	1:50-1:200

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Validation Data

Image



Description

Immunofluorescence staining of HeLa cells with PACO63519 at 1:50, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).