

## HECW1 Antibody

PACO63991

### Description

---

This HECW1 Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

### Product Information

---

<b>SKU:</b>	PACO63991
<b>Contents:</b>	50µl Bradford Reagent: 1 vial (2ml)
<b>Category:</b>	-
<b>Synonyms:</b>	C2 and WW domain-containing protein 1 antibody, E3 ubiquitin-protein ligase HECW1 antibody, HECT antibody, HECT type E3 ubiquitin ligase antibody, HECT, C2 and WW domain containing E3 ubiquitin protein ligase 1 antibody, HECT, C2 and WW domain-containing protein 1 antibody, HECW1 antibody, Hecw1 antibody, HECW1_HUMAN antibody, hNEDL1 antibody, NEDD4 like ubiquitin protein ligase 1 antibody, NEDD4-like E3 ubiquitin-protein ligase 1 antibody, NEDL1 antibody
<b>Clone:</b>	Polyclonal
<b>Applications:</b>	<b>ELISA</b> <b>IF</b>
<b>Conjugation:</b>	Non-conjugated
<b>Reactivity:</b>	Human

### Antibody Data

---

<b>Isotype:</b>	IgG
<b>Uniprot:</b>	Q76N89
<b>Host Species:</b>	Rabbit
<b>Purification:</b>	>95%, Antigen Affinity purified & Affinity purified
<b>Immunogen:</b>	Recombinant Human E3 ubiquitin-protein ligase HECW1 protein (751-900AA)
<b>Immunogen Species:</b>	Homo sapiens (Human)

**Buffer:** Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

**Form:** Liquid

## Preparation & Storage

---

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  
Store Bradford Reagent at Room Temperature for 1 Year.


<b>Recommended Dilutions:</b>	<b>Application</b>	<b>Recommended Dilution</b>
	IF	1:50-1:200

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

## Validation Data

---

**Image**



**Description**

Immunofluorescence staining of A549 cells with PACO63991 at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).