Recombinant EpCAM/TROP-1/TACSTD1 Monoclonal Antibody

Description

This Recombinant EpCAM/TROP-1/TACSTD1 Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	RMEA0149
Contents:	Recombinant EpCAM/TROP-1/TACSTD1 Monoclonal Antibody (50 µL, 100 µL) Bradford Reagent: 1 vial (2ml)
Clone:	C110135
Applications:	WB, IHC-P, ICC/IF, IP
Reactivity:	Human, Mouse, Rat


Antibody Data

Uniprot ID:	P16422
Size:	50 µL, 100 µL
Host Species:	Rabbit
Isotype:	Rabbit IgG
Conjugation:	Unconjugated
Buffer:	0.2 µm filtered solution in PBS


Preparation & Storage

Storage:	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Store Bradford Reagent at Room Temperature for 1 Year.
Shipping:	Gel Pack
Preparation:	WB: 1:1000 ; IHC-P: 1:1000-1:10000 ; ICC/IF: 1:100 ; IP: 1:20-1:50
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol
Notes:	Centrifuge before opening to ensure complete recovery of vial contents.


Validation Data




Immunofluorescent analysis of MCF-7 cells (green), and DAPI (blue).



Immunohistochemical analysis of paraffin-embedded Human colon cancer tissue sections using EpCAM/TROP-1/TACSTD1 Antibody (RMEA0149) at a dilution of 1:1500 (10× lens). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0) for 20 min prior to IHC staining.



Immunohistochemical analysis of paraffin-embedded Mouse colon tissue sections using EpCAM/TROP-1/TACSTD1 Antibody (RMEA0149) at a dilution of 1:1500 (10× lens). Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0) for 20 min prior to IHC staining.



Immunohistochemical analysis of paraffin-embedded Rat colon tissue sections using EpCAM/TROP-1/TACSTD1 Antibody (RMEA0149) at a dilution of 1:1500 (10× lens). Heat mediated antigen retrieval was performed with Tris- EDTA buffer (pH 9.0) for 20 min prior to IHC staining.

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.