Recombinant Stathmin Monoclonal Antibody

Description

This Recombinant Stathmin Monoclonal Antibody is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	RMEA1288
Contents:	Recombinant Stathmin Monoclonal Antibody (50 µL, 100 µL) Bradford Reagent: 1 vial (2ml)
Clone:	C111288
Applications:	WB, IHC, IF, IP
Reactivity:	Human, Mouse, Rat


Antibody Data

Uniprot ID:	P16949
Size:	50 µL, 100 µL
Host Species:	Rabbit
Isotype:	Rabbit IgG
Conjugation:	Unconjugated
Buffer:	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Preparation & Storage

Storage:	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Store Bradford Reagent at Room Temperature for 1 Year.
Shipping:	Gel Pack
Preparation:	WB 1:500-1:2000, IHC 1:200-1:1000, IF 1:50, IP 1:25-1:50
Protein Quantification (Optional):	To quantify total protein levels, use the Bradford Reagent included in this kit. Visit https://www.assaygenie.com/bradford-protein-assay-protocol/ to view the full protocol
Notes:	Centrifuge before opening to ensure complete recovery of vial contents.


Validation Data




Immunofluorescent analysis of (4% Paraformaldehyde) Jurkat cells using anti-Stathmin Monoclonal Antibody at dilution of 1:50.




Immunohistochemistry of paraffin-embedded Human lung squamous carcinoma using Stathmin Monoclonal Antibody at dilution of 1:1000.




Immunohistochemistry of paraffin-embedded Human tonsil using Stathmin Monoclonal Antibody at dilution of 1:1000.




Immunohistochemistry of paraffin-embedded Mouse cerebrum using Stathmin Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded Rat cerebrum using Stathmin Monoclonal Antibody at dilutionfixed of 1:1000.



Immunoprecipitation analysis using anti-Stathmin Monoclonal Antibody. Western blot was performed from the immunoprecipitate using Stathmin Monoclonal Antibody at a dilution of 1:50. Lane 1: 10% Input, Lane 2: Stathmin Monoclonal Antibody, Lane 3: Rabbit monoclonal IgG Isotype Observed-MW:18 kDa Calculated-MW:17 kDa



Western Blot with Stathmin Monoclonal Antibody at dilution of 1:2000. Lane 1: Jurkat, Lane 2: Neuro-2a, Lane 3: Mouse brain Observed-MW:18 kDa Calculated-MW:17 kDa

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.