# **Nanodisc Human CLCN3-Strep Protein**



## **HDFP1243**

### **Product Information**

Product SKU: HDFP1243 Expression Host: HEK293 Size: 10μg

**Target**: CLCN3 **Tag**: C-Flag&Strep Tag

#### **Additional Information**

**Conjugate**: Unconjugated **Uniprot ID**: P51790

Molecular Weight: The human full length CLCN3-Strep protein has a MW of 91 kDa

#### **Protein Information**

**Background**: This gene encodes a member of the voltage-gated chloride channel (CIC) family. The

encoded protein is present in all cell types and localized in plasma membranes and

in intracellular vesicles. It is a multi-pass membrane protein which contains a CIC

domain and two additional C-terminal CBS (cystathionine beta-synthase) domains.

The CIC domain catalyzes the selective flow of CI- ions across cell membranes, and

the CBS domain may have a regulatory function. This protein plays a role in both

acidification and transmitter loading of GABAergic synaptic vesicles, and in smooth

muscle cell activation and neointima formation. This protein is required for

lysophosphatidic acid (LPA)-activated CI- current activity and fibroblast-to-

myofibroblast differentiation. The protein activity is regulated by Ca(2 )/calmodulin-

dependent protein kinase II (CaMKII) in glioma cells. Multiple alternatively spliced

transcript variants encoding different isoforms have been identified. [provided by

RefSeq, Aug 2011]

**Synonyms**: CLC3, ClC-3

**Protein Description**: Human CLCN3-Strep full length protein-synthetic nanodisc

Formulation: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH

8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please

see Certificate of Analysis for specific instructions. Do not use solvents with a pH

below 6.5 or those containing high concentrations of divalent metal ions (greater

than 5 mM) in subsequent experiments.

Protein Pathways: -

**Protein Families:** Ion Channels: Other.

**Usage**: Research use only

**Storage & Shipping**: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing

and thawing). Lyophilized proteins are shipped at ambient temperature.