## Nanodisc Human FXYD5-Strep Protein



## **HDFP1308**

## **Product Information**

<b>Product SKU</b> :	HDFP1308	Expression Host:	HEK293		Size:	10µg	
Target:	FXYD5	Tag:	C-Flag&St	rep Tag			
Additional Infor	mation						
<b>Conjugate</b> :	Unconjugat	ed Unip	orot ID:	Q96DB9			
Molecular Wei	ght: The human	The human full length FXYD5-Strep protein has a MW of 19.5 kDa					

## **Protein Information**

Background: This gene encodes a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7 invariant and 6 highly conserved amino acids. The approved human gene nomenclature for the family is FXYD-domain containing ion transport regulator. Mouse FXYD5 has been termed RIC (Related to Ion Channel). FXYD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. Transmembrane topology has been established for two family members (FXYD1 and FXYD2), with the N-terminus extracellular and the C-terminus on the cytoplasmic side of the membrane. This gene product, FXYD5, is a glycoprotein that functions in the up-regulation of chemokine production, and it is involved in the reduction of cell adhesion via its ability to down-regulate E-cadherin. It also promotes metastasis, and has been linked to a variety of cancers. Alternative splicing results in multiple transcript variants. [RefSeq curation Kathleen J. Sweadner, by Ph.D., sweadner@helix.mgh.harvard.edu., Sep 2009] Synonyms: DYSAD, HSPC113, IWU1, KCT1, OIT2, PRO6241, RIC Human FXYD5-Strep full length protein-synthetic nanodisc **Protein Description:** 

Formulation:	tion: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH		
	8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please		
	see Certificate of Analysis for specific instructions. Do not use solvents with a pH		
	below 6.5 or those containing high concentrations of divalent metal ions (greater		
	than 5 mM) in subsequent experiments.		
Protein Pathways:	-		
Protein Families:	Ion Channels: Other.		
Usage:	Research use only		
Storage & Shipping:	ing: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not		
	intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing		
	and thawing). Lyophilized proteins are shipped at ambient temperature.		