

## Recombinant Human CD69 Protein

RPCB0154

### Protein Information

---

<b>Size:</b>	10 µg , 20 µg , 50 µg , 100 µg	<b>Tag:</b>	C-His
<b>Reactivity:</b>	Human	<b>Expressed Host:</b>	HEK293 cells
<b>Calculated MW:</b>	16.81 kDa	<b>Observed MW:</b>	20-26 kDa

### Background

---

Early activation antigen CD69, also known as activation inducer molecule (AIM), is a single-pass type II membrane protein. Recently, cDNA clones encoding human and mouse CD69 were isolated and showed CD69 to be a member of the C-type lectin superfamily. It is one of the earliest cell surface antigens expressed by T cells following activation. Once expressed, CD69 acts as a costimulatory molecule for T cell activation and proliferation. In addition to mature T cells, CD69 is inducibly expressed by immature thymocytes, B cells, natural killer (NK) cells, monocytes, neutrophils and eosinophils, and is constitutively expressed by mature thymocytes and platelets. CD69 is involved in lymphocyte proliferation and functions as a signal transmitting receptor in lymphocytes, natural killer (NK) cells, and platelets. The structure, chromosomal localization, expression and function of CD69 suggest that it is likely a pleiotropic immune regulator, potentially important in the activation and differentiation of a wide variety of hematopoietic cells. This membrane molecule transiently expresses on activated lymphocytes, and its selective expression in inflammatory infiltrates suggests that it plays a role in the pathogenesis of inflammatory diseases. CD69 plays a crucial role in the pathogenesis of allergen-induced eosinophilic airway inflammation and hyperresponsiveness and that CD69 could be a possible therapeutic target for asthmatic patients.

### Properties


---

<b>Synonyms:</b>	CD69, AIM, BL-AC/P26, CLEC2C, EA1, GP32/28, MLR-3
<b>Gene ID:</b>	969
<b>Endotoxin:</b>	< 0.1 EU/µg of the protein by LAL method.
<b>Description:</b>	High quality, high purity and low endotoxin recombinant Recombinant Human CD69 Protein (RPCB0154), tested reactivity in HEK293 cells and has been validated in SDS-PAGE. 100% guaranteed.
<b>Purity:</b>	≥ 95 % as determined by SDS-PAGE.
<b>Storage:</b>	Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.



## Validation Data

---



Recombinant Human CD69 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.