

RPCB0895

Product Information

Product SKU:	RPCB0895	Gene ID:	1634	Size:	10µg
Tag:	C-His	Reactivity:	Human		

Additional Information

Expression Host:	HEK293 cells	Swissprot:	P07585-1
Purity:	> 95% by SDS-PAGE.		

Protein Information

Background: Decorin is a ubiquitous small cellular or pericellular matrix proteoglycan and is closely related in structure to biglycan protein. It belongs to the small leucine-rich proteoglycan (SLRP) family and consists of a core protein and a covalently linked glycosaminoglycan chain which is either chondroitin sulfate (CS) or dermatan sulfate (DS). As a component of connective tissue, decorin interacts with several extracellular matrix components, such as type I collagen and fibronectin, and plays a role in matrix assembly. Decorin resides in the tumor microenvironment and affects the biology of various types of cancer by downregulating the activity of several receptors involved in cell growth and survival. Decorin binds to and modulates the signaling of the epidermal growth factor receptor and other members of the ErbB family of receptor tyrosine kinases. It exerts its antitumor activity by a dual mechanism: via inhibition of these key receptors through their physical downregulation coupled with attenuation of their signaling, and by binding to and sequestering TGFbeta. Decorin also modulates the insulin-like growth factor receptor and the low-density lipoprotein receptor-related protein 1, which indirectly affects the TGFbeta receptor pathway. Decorin plays significant roles in tissue development and assembly, as well as playing both direct and indirect signaling roles.

- Protein Description:** High quality, high purity and low endotoxin recombinant Recombinant Human Decorin/PG-S2/DCN Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.
- Endotoxin:** Please contact us for more information.
- Formulation:** Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.
- Storage:** Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.