

RPCB1227

---

**Product Information**

<b>Product SKU:</b>	RPCB1227	<b>Gene ID:</b>	780	<b>Size:</b>	10µg
<b>Tag:</b>	C-His	<b>Reactivity:</b>	Human		

---

**Additional Information**

<b>Expression Host:</b>	-	<b>Swissprot:</b>	Q08345-1
<b>Purity:</b>	> 95% by SDS-PAGE.		

---

**Protein Information**

**Background:** Discoidin domain receptor family, member 1 (DDR1), also known as or CD167a (cluster of differentiation 167a), and Mammary carcinoma kinase 10 (MCK10), belongs to a subfamily of tyrosine kinase receptors with an extracellular domain homologous to Dictyostellium discoideum protein discoidin 1. Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases are involved in the regulation of cell growth, differentiation and metabolism. Expression of DDR1/MCK10/CD167 is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. DDR1/MCK10/CD167 plays an important role in regulating attachment to collagen, chemotaxis, proliferation, and MMP production in smooth muscle cells. DDR1 functions in a feedforward loop to increase p53 levels and at least some of its effectors. Inhibition of DDR1 function resulted in strikingly increased apoptosis of wild-type p53-containing cells in response to genotoxic stress through a caspase-dependent pathway.

**Protein Description:** High quality, high purity and low endotoxin recombinant Recombinant Human HGK2/DDR1/CD167a Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.

**Endotoxin:** <0.1EU/µg

**Formulation:** Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

**Storage:** Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.