

## RPCB1259

---

**Product Information**

<b>Product SKU:</b>	RPCB1259	<b>Gene ID:</b>	2335	<b>Size:</b>	10µg
<b>Tag:</b>	C-His	<b>Reactivity:</b>	Human		

---

**Additional Information**

<b>Expression Host:</b>	HEK293 cells	<b>Swissprot:</b>	P02751
<b>Purity:</b>	> 90% by SDS-PAGE.		

---

**Protein Information**

**Background:** Fibronectin (FN) is a glycoprotein component of the extracellular matrix of the extracellular matrix (ECM) with roles in embryogenesis, development, and wound healing. More recently, FN has emerged as player in platelet thrombus formation and diseases associated with thrombosis including vascular remodeling, atherosclerosis, and cardiac repair following a myocardial infarct. Each monomer of FN consists of three types of homologous repeating units, that is 12 type I repeats, two type II repeats and 15-17 type III repeats. The occurrence of multiple isoforms results from alternative mRNA splicing of the ED-A, ED-B and III-CS regions, and subsequent post-translational modification. As an ECM component and one of the primary cell adhesion molecules, Fibronectin can be a ligand for fibrin, heparin, chondroitin sulfate, collagen/gelatin, as well as many integrin receptors through which FN mediates the variety of cellular signaling pathways. The study of solid human tumors showed among the early signs of malignant transformation the fragmentation of pericellular FN, concomitant with the increase of its production by the peritumoral stroma. These results should encourage further investigations concerning the potential importance of Fn production and breakdown during cancer progression. FN1 expression has been described to increase significantly from the morula towards the early blastocyst stage, suggesting that FN1 may also be involved in early blastocyst formation. The fragment 2 of FN comprises the first 7 FN type III repeats

and is suggested to be important for self association during fibril growth via the key module III2.

- Protein Description:** High quality, high purity and low endotoxin recombinant Recombinant Human Fibronectin/CIG/FN1 Protein , tested reactivity in HEK293 cells and has been validated in SDS-PAGE.100% guaranteed.
- Endotoxin:** < 0.1 EU/μg of the protein by LAL method.
- Formulation:** Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.
- Storage:** Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.