

Recombinant Mouse AGO2/Argonaute 2/EIF2C2 Protein (His Tag)

RPES0312

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES0312

Purity: > 80 % as determined by reducing SDS-PAGE.

Contents: 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Mouse

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: 1110029L17Rik,
2310051F07Rik, AI225898, AL022874,
AW546247, ENSMUSG00000072493,
Eif2c2, Gerp95, Gm10365, mKIAA4215

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: Baculovirus-Insect Cells

Bio-Activity: Not validated for activity

Calculated MW: 99.0 kDa

Formulation: Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% Glycerol pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 105 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: Q8CJG0

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: Baculovirus-Insect Cells-derived Mouse AGO2/Argonaute 2/EIF2C2 protein Met 1- Ala 860, with an N-terminal His

Sequence: Met 1-Ala 860

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.