

Recombinant Human BCL2/Bcl-2 Protein (His Tag)

RPES0599

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES0599

Purity: > 90 % as determined by reducing SDS-PAGE.

Contents: 100µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: Please contact us for more information.

Synonyms: Apoptosis Regulator Bcl-2,
Apoptosis regulator Bcl-2, B-cell
Lymphoma 2, BCL2, PPP1R50

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Immobilized human BCL2-His at 10 µg/ml (100 µl/well) can bind biotinylated mouse BCL2L1-His, The EC50 of biotinylated mouse BCL2L1-His is 0.07-0.15 µg/ml.

Calculated MW: 24.7 kDa

Formulation: Lyophilized from sterile 50mM Tris, 20% glycerol, 100mM Arg, pH 8.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 32 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: P10415-1

Source: E.coli-derived Human BCL2/Bcl-2 protein Met 1-Asp 211, with an C-terminal His

Sequence: Met 1-Asp 211

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.