

Recombinant Human Oncostatin M/OSM Protein (His Tag)

RPES0689

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES0689

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 0.01 EU per µg of the protein as determined by the LAL method.

Synonyms: OSM, Oncostatin-M

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Measured by the dose-dependent stimulation of TF-1 cells. The ED50 for this effect is 0.2-1 ng/ml.

Calculated MW: 24.4 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of 20mMTris-HCl, 1mM EDTA, 200mM NaCl, pH 7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 28 kDa

Reconstitution: Please refer to the printed manual for detailed information. Measured by the dose-

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

dependent stimulation of TF-1 cells. The ED50 for this effect is 0.2-1 ng/ml.

Accession: P13725

Source: E.coli-derived Human Oncostatin M/OSM protein Ala26-Arg221, with an N-terminal His

Sequence: Ala26-Arg221

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.