

Recombinant Human CXCL7/NAP-2 Protein

RPES1406

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES1406

Purity: > 98 % as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 0.1 EU per µg of the protein as determined by the LAL method.

Synonyms: C-X-C Motif Chemokine 7, CTAP3, CXCL7, LDGF, Leukocyte-Derived Growth Factor, MDGFSmall-Inducible Cytokine B7, Macrophage-Derived Growth Factor, PBP, PPBP, Platelet Basic Protein, SCYB7, TGB1, THBGB1

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Measure by its ability to chemoattract BaF3 cells transfected with human CXCR2. The ED50 for this effect is <0.5 ng/mL.

Calculated MW: 8.4 kDa

Formulation: Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 11 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: P02775

Source: E.coli-derived Human CXCL7, NAP-2 protein Ala59-Asp128, with an N-terminal His

Sequence: Ala59-Asp128

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.