

Recombinant Human HSP40/DNAJB1 Protein (His Tag)

RPES2501

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES2501

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: DNAJ1, DNAJB1, DnaJ Homolog Subfamily B Member 1, DnaJ Protein Homolog 1, HDJ1, HSP40, HSPF1, Heat Shock 40 kDa Protein 1, Heat Shock Protein 40, Human DnaJ Protein 1, hDj-1

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Not validated for activity

Calculated MW: 39.1 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, 1mM EDTA, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 38 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: P25685

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: E.coli-derived Human HSP40, DNAJB1 protein Gly2-Ile340, with an C-terminal His

Sequence: Gly2-Ile340

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.