

Recombinant Cynomolgus CD3e/CD3ε Protein (Fc Tag)

RPES2899

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES2899

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Cynomolgus Macaques

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: CD3 epsilon, CD3-epsilon, CD3e, CD3e antigen, CD3e molecule, FLJ18683, T-cell antigen receptor complex, T-cell surfa, T-cell surface antigen T3/Leu-4 epsilon chain, T3E, epsilon (CD3-TCR complex), epsilon polypeptide (TiT3 complex), epsilon subunit of T3

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-Fc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 38 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of 50 mM Tris-HCl, 100 mM Glycine, pH 7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 38-55 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: Q95LI5

Source: HEK293 Cells-derived
Cynomolgus macaques CD3e/CD3ε
protein Gln22-Asp117, with an C-terminal
Fc

Sequence: Gln22-Asp117

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.