

Recombinant Human Ephrin-A1/EFNA1 Protein (His Tag)

RPES2966

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES2966

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: EFNA1, EPH-Related Receptor Tyrosine Kinase Ligand 1, EPLG1, Ephrin-A1, Immediate Early Response Protein B61, LERK-1, LERK1, TNF Alpha-Induced Protein 4, TNFAIP4, Tumor Necrosis Factor Alpha-Induced Protein 4

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 20.4 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 4% Trehalose, 4% Mannitol, 0.02% Tween 80, pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 24-28 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: AAH32698.1

Source: HEK293 Cells-derived Human Ephrin-A1, EFNA1 protein Asp19-Ser182, with an C- terminal His

Sequence: Asp19-Ser182

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.