

Recombinant Human IL13RA2/CD213A2 Protein (His &Fc Tag)

RPES3990

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES3990

Purity: > 75 % as determined by reducing SDS-PAGE.

Contents: 100µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: IL13RA, CD213A, IL13RA2, CD213A2, CT19, IL-13R, IL13BP, IL13R, IL-13Ra2

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His-Fc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Immobilized recombinant human IL13RA2 at 8 µg/ml (100 µl/well) can bind IL13 with a linear range of 0.25-8.0 ng/ml.

Calculated MW: 65.0 kDa

Formulation: Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 90-100 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: NP_000631.1

Source: HEK293 Cells-derived Human IL13RA2/CD213A2 protein Met 1-Leu 342, with an C- terminal His & Fc

Sequence: Met 1-Leu 342

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.