

# Recombinant Human HMGB1/HMG1 Protein (Fc Tag)

RPES4393

## Description

---

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Protein Information

---

**SKU:** RPES4393

**Purity:** > 97 % as determined by reducing SDS-PAGE.

**Contents:** 50µg  
Bradford Reagent: 1 vial (2ml)

**Concentration:** -

**Species:** Human

**Endotoxin:** < 1.0 EU per µg of the protein as determined by the LAL method.

**Synonyms:** HMG-1, HMG1, HMG3, SBP-1

**Storage:** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.  
Store Bradford Reagent at Room Temperature for 1 year.

**Tag:** N-hFc

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Expression Host:** HEK293 Cells

**Bio-Activity:** Immobilized recombinant mouse AGER at 2 µg/ml (100 µl/well) can bind human HMGB1. The EC50 of human HMGB1 is 0.23 µg/ml.

**Calculated MW:** 51.5 kDa

**Formulation:** Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

**Observed MW:** 55-60 kDa

**Reconstitution:** Please refer to the printed manual for detailed information.

**Manufacturers Statement:** This final kit system is assembled and quality-released by Assay Genie Limited.

**Accession:** NP\_002119.1

**Source:** HEK293 Cells-derived Human HMGB1/HMG1 protein Met 1-Glu 215, with an N- terminal hFc

**Sequence:** Met 1-Glu 215

**Form:** Lyophilized powder

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.