

Recombinant Mouse Adiponectin/AdipoQ Protein (E.coli, His Tag)

RPES4743

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES4743

Purity: > 90 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Mouse

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: 30 kDa Adipocyte Complement-Related Protein, ACRP30, ADIPOQ, Adipocyte, Adipocyte complement-related 30 kDa protein, Adiponectin, Adipose Most Abundant Gene Transcript 1 Protein, C1q and Collagen Domain-Containing Protein, Gelatin-Binding Protein, apM-1

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Not validated for activity

Calculated MW: 27.2 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 30 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: Q60994

Source: E.coli-derived Mouse Adiponectin, AdipoQ protein Glu18-Asn247 , with an N-terminal His

Sequence: Glu18-Asn247

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.