

Recombinant Mouse PARP-1 Protein (His Tag)

RPES4756

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES4756

Purity: > 85 % as determined by reducing SDS-PAGE.

Contents: 100µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Mouse

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: 5830444G22Rik, A1893648, ARTD1, Adprp, Adprt1, C80510, PARP, PPOL, parp-1, sPARP-1

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: Baculovirus-Insect Cells

Bio-Activity: Immobilized mouse PARP1 at 10 µg/mL (100 µl/well) can bind biotinylated human HSP70, The EC50 of biotinylated human HSP70 is 0.021 µg/mL.

Calculated MW: 115 kDa

Formulation: Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol, 0.1mM TCEP Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 75 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: NP_031441.2

Source: Baculovirus-Insect Cells-derived
Mouse PARP-1 protein Met 1-Trp 1014,
with an N- terminal His

Sequence: Met 1-Trp 1014

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.