

Recombinant Mouse CD16-2/FCGR4 Protein (His&AVI Tag), Biotinylated RPES5101

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES5101

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Mouse

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: 4833442P21Rik Protein, CD16-2 Protein, FcgRIV Protein, FcγRIV Protein, FcγR3a Protein, FcγR3 Protein, Mouse

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His-Avi

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: 1. Measured by its ability to bind human IgG1 (Fc) in a functional ELISA. 2. Labeling ratio of biotin to protein: 0.6

Calculated MW: 24.2 kDa

Formulation: Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: -

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: NP_653142.2

Source: HEK293 Cells-derived Mouse CD16-2/FCGR4 protein Met 1-Gln 203, with an C- terminal His & Avi

Sequence: Met 1-Gln 203

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.