

Recombinant Human CD40/TNFRSF5 Protein (Fc Tag)

RPES5182

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES5182

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: B-Cell Surface Antigen CD40, Bp50, CD40, CD40L Receptor, CDw40, TNFRSF5, Tumor Necrosis Factor Receptor Superfamily member 5

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-Fc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Immobilized Human CD40L-His-Flag at 10µg/ml (100 µl/well) can bind Human CD40- Fc. The ED50 of Human CD40-Fc is 0.45 µg/ml.

Calculated MW: 46.3 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 50-65 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: P25942

Source: HEK293 Cells-derived Human CD40, TNFRSF5 protein Glu21-Arg193, with an C- terminal Fc

Sequence: Glu21-Arg193

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.