

# Recombinant Human CD30L/TNFSF8 Protein (His Tag)

RPES5995

## Description

---

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Protein Information

---

**SKU:** RPES5995

**Purity:** > 90% as determined by reducing SDS-PAGE.

**Contents:** 100µg, 20µg, 500µg, 1mg  
Bradford Reagent: 1 vial (2ml)

**Concentration:** -

**Species:** Human

**Endotoxin:** < 1.0 EU/mg of the protein as determined by the LAL method

**Synonyms:** CD antigen CD153, CD30 ligand, CD30-L, CD30L, CD30LG, TNFSF8, Tumor necrosis factor ligand superfamily member 8

**Storage:** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.  
Store Bradford Reagent at Room Temperature for 1 year.

**Tag:** C-His

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Expression Host:** HEK293 Cells

**Bio-Activity:** Not validated for activity

**Calculated MW:** 18.8 kDa

**Formulation:** Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

**Observed MW:** 30-40 kDa

**Reconstitution:** It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

**Accession:** P32971

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Manufacturers Statement:** This final kit system is assembled and quality-released by Assay Genie Limited.

**Source:** HEK293 Cells-derived Human CD30L/TNFSF8 protein Gln63-Asp234, with an C- terminal His

**Sequence:** Gln63-Asp234

**Form:** Lyophilized powder

**Notes:** Centrifuge before opening to ensure complete recovery of vial contents.