

Recombinant Human IL-8/CXCL8 Protein (aa 28-99, Fc Tag)

RPES6165

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6165

Purity: > 90 % as determined by reducing SDS-PAGE.

Contents: 100µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: CXCL, Interleukin, C-X-C motif chemokine, Chemokine (C-X-C motif) ligand, IL8/CXCL, GCP, NAP, CXCL8, GCP-1, GCP1, IL8, LECT, LUCT, LYNAP, MDNCF, MONAP, NAF, NAP-1, NAP1, 3-10C, AMCF-I, IL8/CXCL8, K60, Interleukin-8, IL-8, Emoctakin, C-X-C motif chemokine 8, Chemokine (C-X-C motif) ligand 8, Granulocyte chemotactic protein 1 (GCP-1), Monocyte-derived neutrophil chemotactic factor (MDNCF), (Ala-IL-8)77, (Ser-IL-8)72, 9E3, Beta thromboglobulin like protein, CEF-4, chemokine, CXC motif, GCP/IL-8 protein I, GCP/IL-8 protein II, GCP/IL-8 protein III, GCP/IL-8 protein IV, GCP/IL-8 protein V, GCP/IL-8 protein VI, Granulocyte chemotactic protein 1, IL-8(1-77), IL-8(9-77), IL8/NAP1 form I, IL8/NAP1 form II, IL8/NAP1 form III, IL8/NAP1 form IV, IL8/NAP1 form V, IL8/NAP1 form VI, Inteleukin 8, ligand 8, Lymphocyte-derived neutrophil-activating factor, MDNCF-b, MDNCF-c, Monocyte-derived neutrophil chemotactic factor, Monocyte-derived neutrophil-activating peptide, Neutrophil activating peptide 1, Neutrophil activating protein 1, Neutrophil-activating factor, Neutrophil-activating protein 1, Protein 3

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

10C, Protein 3-10C, SCYB8, Small inducible cytokine subfamily B member 8, T cell chemotactic factor, T-cell chemotactic factor, TSG1

Tag: N-hFc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 35.0 kDa

Formulation: Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 40 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: NP_000575.1

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Source: HEK293 Cells-derived Human IL-8/CXCL8 protein Ser 28-Ser 99, with an N-terminal hFc

Sequence: Ser 28-Ser 99

Notes: Centrifuge before opening to ensure complete recovery of vial contents.

Form: Lyophilized powder

