

Recombinant Human IL-23 alpha & Mouse IL-12 beta Heterodimer (C-RPES6211

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6211

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human, Mouse

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: CLMF p40, IL-12 subunit p40, IL-23p19, NKSF2, SGRF

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Measured by its ability to induce STAT reporter activity in 293F human embryonic kidney cells. The ED50 for this effect is 357.68 ng/ml.

Calculated MW: 56.9 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 65-80 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: Q9NPF7&P43432

Source: HEK293 Cells-derived Human & Mouse IL-23A & IL-12B protein Arg20-Pro189& Met23-Ser335, with an C-terminal His

Sequence: Arg20-Pro189&Met23-Ser335

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.