

Recombinant Human P-selectin/CD62P (C-Fc)

RPES6373

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6373

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: CD62 antigen-like family member P, GMP-140, Granule membrane protein 140, LECAM3, Leukocyte-endothelial cell adhesion molecule 3, P-selectin, PADGEM

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-Fc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 106.9 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 130-160 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: P16109

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: HEK293 Cells-derived Human P-selectin, CD62P protein Trp42-Ala771, with an C- terminal Fc

Sequence: Trp42-Ala771

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.