

Recombinant Human IL-5 protein(His Tag)

RPES6409

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6409

Purity: > 98 % as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 0.1 EU per µg of the protein as determined by the LAL method.

Synonyms: nterleukin, Interleukin, IL5, EDF, IL-5, TRF, BCDFII, B-cell differentiation factor I, B-cell growth factor I, BCGF-II, Cytotoxic T-lymphocyte inducer, E-CSF, Eosinophil differentiation factor, ESGF, Interleukin-5, nterleukin-5, T-cell replacing factor

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Measure by its ability to induce TF-1 cells proliferation. The ED50 for this effect is <0.3 ng/mL.

Calculated MW: 14.1 kDa

Formulation: Lyophilized from sterile PBS,pH 8.0. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 14 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: P05113

Source: E.coli-derived Human IL-5 protein
Ile 20-Ser 134, with an C-terminal His

Sequence: Ile 20-Ser 134

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.