

Recombinant Human TGF beta 3 protein(His Tag)

RPES6506

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6506

Purity: > 98 % as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 0.1 EU per µg of the protein as determined by the LAL method.

Synonyms: ARVD, ARVD1, LDS5, RNHF, TGF-B3, TGFB3

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Measure by its ability to inhibit IL-4-induce proliferation in HT-2 cells. The ED50 for this effect is <50 pg/mL. The specific activity of recombinant human TGF beta 3 is > 2 x 10⁷ IU/mg.

Calculated MW: 13.7 kDa

Formulation: Lyophilized from sterile 20 mM sodium citrate, 0.2 M NaCl, pH 3.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 13 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Accession: P10600

Source: E.coli-derived Human TGF beta 3 protein Ala 301-Ser 412, with an C-terminal His

Sequence: Ala 301-Ser 412

Form: Lyophilized powder

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.