

Recombinant Rhesus Macaque Angiopoietin-2/ANG2 (C-Fc)

RPES6789

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES6789

Purity: > 95 % as determined by reducing SDS-PAGE.

Contents: 50µg, 10µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Rhesus Macaque

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: AGPT2, ANG-2, ANG2, ANGPT2, Angiopoietin-2, Tie2-ligand, angiopoietin 2, angiopoietin-2B, angiopoietin-2a, angiopoitin 2

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-Fc

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 81.7 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: 90-120 kDa

Reconstitution: Please refer to the printed manual for detailed information.

Accession: XM_001097949

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: HEK293 Cells-derived Rhesus Macaque Angiotensin-2/ANG2 protein Tyr19-Phe495, with an C-terminal Fc

Sequence: Tyr19-Phe495

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.