

Recombinant SARS-CoV-2 Spike S1+S2 ECD

RPES7085

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES7085

Purity: > 90 % as determined by reducing SDS-PAGE.

Contents: 20µg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: SARS-CoV-2

Endotoxin: < 1.0 EU per µg of the protein as determined by the LAL method.

Synonyms: SARS-CoV-, coronavirus s1 Protein, coronavirus s2 Protein

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: Baculovirus-Insect Cells

Bio-Activity: Not validated for activity

Calculated MW: 134 kDa

Formulation: Lyophilized from sterile PBS (contains 300 mM NaCl), pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Observed MW: -

Reconstitution: Please refer to the printed manual for detailed information.

Accession: YP_009724390.1

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: Baculovirus-Insect Cells-derived SARS-CoV-2 SARS-CoV-2 Spike S1+S2 ECD (L18F, D80A, D215G, ΔLAL242-244, R246I, K417N, E484K, N501Y, D614G, A701V) protein Met1- Pro1213, with an C-terminal His

Sequence: Met1-Pro1213

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.