

Recombinant Human GAPDH Protein (His Tag)

RPES8588

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES8588

Purity: > 90% as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg, 500µg, 1mg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 10 EU/mg of the protein as determined by the LAL method

Synonyms: Aging associated gene 9 protein, Epididymis secretory sperm binding protein Li 162eP, G3P, G3PD, G3PDH, GAPD, GAPDH, Glyceraldehyde 3 phosphate dehydrogenase, Glyceraldehyde-3-phosphate dehydrogenase, HEL S 162eP, Peptidyl-cysteine S-nitrosylase GAPDH

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to - 80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.

Store Bradford Reagent at Room Temperature for 1 year.

Tag: N-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: E.coli

Bio-Activity: Not validated for activity

Calculated MW: 37.1 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Observed MW: 34-37 kDa

Reconstitution: It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Accession: P04406

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: E.coli-derived Human GAPDH protein Met1-Glu333, with an N-terminal His

Sequence: Met1-Glu333

Form: Lyophilized powder

this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Notes: Centrifuge before opening to ensure complete recovery of vial contents.