

# Recombinant Human IL-17F Protein (His Tag)

RPES8776

## Description

---

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Protein Information

---

**SKU:** RPES8776

**Purity:** > 95% as determined by reducing SDS-PAGE.

**Contents:** 100µg, 20µg, 500µg, 1mg  
Bradford Reagent: 1 vial (2ml)

**Concentration:** -

**Species:** Human

**Endotoxin:** < 10 EU/mg of the protein as determined by the LAL method

**Synonyms:** Cytokine ML, Interleukin, CANDF, IL17F, CANDF6, IL-17F, ML-1, ML1, Cytokine ML-1, IL24, IL-24, Interleukin-17F, Interleukin-24

**Storage:** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to - 80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.  
Store Bradford Reagent at Room Temperature for 1 year.

**Tag:** C-His

**Shipping:** This product is provided as lyophilized powder which is shipped with ice packs.

**Expression Host:** E.coli

**Bio-Activity:** Not validated for activity

**Calculated MW:** 14.5 kDa

**Formulation:** Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

**Observed MW:** 16 kDa

**Reconstitution:** It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

**Accession:** Q96PD4

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

**Manufacturers Statement:** This final kit system is assembled and quality-released by Assay Genie Limited.

**Source:** E.coli-derived Human IL-17F  
protein Arg31-Gln163, with an C-terminal  
His

**Sequence:** Arg31-Gln163

**Form:** Lyophilized powder

**Notes:** Centrifuge before opening to ensure complete  
recovery of vial contents.