

Recombinant Human SIGLEC2/CD22 Protein (His Tag)

RPES9228

Description

This high-purity recombinant protein is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Protein Information

SKU: RPES9228

Purity: > 95% as determined by reducing SDS-PAGE.

Contents: 100µg, 20µg, 500µg, 1mg
Bradford Reagent: 1 vial (2ml)

Concentration: -

Species: Human

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Synonyms: B-cell receptor CD22, B-lymphocyte cell adhesion molecule, BL-CAM, CD22, CD22 antigenMGC130020, CD22 molecule, SIGLEC-2, SIGLEC2FLJ22814, Siglec-2, T-cell surface antigen Leu-14, sialic acid binding Ig-like lectin 2

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Store Bradford Reagent at Room Temperature for 1 year.

Tag: C-His

Shipping: This product is provided as lyophilized powder which is shipped with ice packs.

Expression Host: HEK293 Cells

Bio-Activity: Not validated for activity

Calculated MW: 76.2 kDa

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Observed MW: 100-140 kDa

Reconstitution: It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Accession: P20273

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

Manufacturers Statement: This final kit system is assembled and quality-released by Assay Genie Limited.

Source: HEK293 Cells-derived Human
SIGLEC2 protein Asp20-Arg687, with an C-
terminal His

Sequence: Asp20-Arg687

Form: Lyophilized powder

Notes: Centrifuge before opening to ensure complete
recovery of vial contents.