

Anti-Mouse Ly-6G/Ly-6C (Gr-1) In Vivo Antibody - Low Endotoxin

IVMB0091

Description

This Anti-Mouse Ly-6G/Ly-6C (Gr-1) In Vivo Antibody - Low Endotoxin is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: IVMB0091

Contents: 1mg, 5mg, 25mg, 50mg, 100mg
Bradford Reagent: 1 vial (2ml)

Synonyms: Gr-1

Category: Monoclonal Antibody

Target: Ly-6G/Ly-6C

Clone: RB6-8C5

Isotype: Rat IgG2b

Applications: 

Specificity: Rat Anti-Mouse GR-1, as known as anti-mouse Ly-6G/Ly-6C (Clone RB6-8C5) has been used to identify peripheral blood neutrophils and deplete granulocytes in vivo. This monoclonal antibody was purified using multi-step affinity chromatography methods such as Protein A or G depending on the species and isotype.

Antibody Data

Reactivity: Mouse

Host species: Rat

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

Expression Host: -

Immunogen: Raised against granulocytes of Mouse origin

Product concentration: ≥ 5.0 mg/ml

Endotoxin Level: < 1.0 EU/mg as determined by the LAL method

Purity: ≥95% Monomer by analytical SEC, >95% by SDS Page

Formulation: This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

Preparation & Storage

Storage: Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at ≤ -70°C. Avoid Repeated Freeze Thaw Cycles.
Store Bradford Reagent at Room Temperature for 1 Year.

Shipping: Next Day 2-8°C

Preparation: Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Recommended Dilution Buffer: In vivo Antibody Diluent pH 7.2

Recommended Usage:	Application	Recommended Usage
	FC	The suggested concentration for this RB6-8C5 antibody for staining cells in flow cytometry is ≤ 0.06 µg per 10 ⁶ cells in a volume of 100 µl or 100 µl of whole blood. Titration of the reagent is recommended for optimal performance for each application.

**Protein
Quantification
(Optional):**

To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol