

Anti-Mouse NK1.1 In Vivo Antibody - Low Endotoxin

IVMB0100

Description

This Anti-Mouse NK1.1 In Vivo Antibody - Low Endotoxin is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

| | |
|----------------------|--|
| SKU: | IVMB0100 |
| Contents: | 1mg, 5mg, 25mg, 50mg, 100mg Bradford Reagent: 1 vial (2ml) |
| Synonyms: | Natural Killer Cells, KR-P1C, NKR-P1B, Ly-55, CD161b, CD161c |
| Category: | Monoclonal Antibody |
| Target: | NK1.1 |
| Clone: | PK136 |
| Isotype: | Mouse IgG2a k |
| Applications: | B CyTOF® Depletion FC In Vivo IP WB |
| Specificity: | This NK1.1 antibody (clone PK136) recognizes mouse NK1.1. |

Antibody Data

| | |
|-------------------------|---|
| Reactivity: | Mouse |
| Host species: | Mouse |
| Expression Host: | - |
| Immunogen: | NK-1+ cells from Mouse spleen and bone marrow |

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

- Product concentration:** ≥ 5.0 mg/ml
- Endotoxin Level:** < 1.0 EU/mg as determined by the LAL method
- Purity:** ≥95% Monomer by analytical SEC, >95% by SDS Page
- Formulation:** This monoclonal NK1.1 antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

Preparation & Storage

Storage: Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at ≤ -70°C. Avoid Repeated Freeze Thaw Cycles.
Store Bradford Reagent at Room Temperature for 1 Year.

Shipping: Next Day 2-8°C

Preparation: Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Recommended Dilution Buffer: In vivo Antibody Diluent pH 7.2

| Recommended Usage: | Application | Recommended Usage |
|---------------------------|--------------------|--|
| | Flow Cytometry | The suggested use of this NK1.1 antibody is ≤ 0.25 µg per 10 ⁶ cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application. |

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol