

Anti-Human CD16 [3G8] In Vivo Antibody - Low Endotoxin

IVMB0112

Description

This Anti-Human CD16 [3G8] In Vivo Antibody - Low Endotoxin is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU: IVMB0112

Contents: 1mg, 5mg, 25mg, 50mg, 100mg
Bradford Reagent: 1 vial (2ml)

Synonyms: FcγRIII, Fc Gamma RIII

Category: Monoclonal Antibody

Target: CD16

Clone: 3G8

Isotype: Mouse IgG1 κ

Applications: **B** **CyTOF®** **FC** **IHC FF** **In Vivo** **IP**

Specificity: Clone 3G8 recognizes an epitope on human CD16. This clone also cross-reacts with non-human primate CD16.

Antibody Data

Reactivity: Baboon, Chimpanzee, Cynomolgus Monkey, Marmoset, Pigtailed Macaque, Rhesus Monkey, Squirrel Monkey, Human

Host species: Mouse

Expression Host: -

Immunogen: Human PMN cells

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

- Product concentration:** ≥ 5.0 mg/ml
- Endotoxin Level:** < 1.0 EU/mg as determined by the LAL method
- Purity:** ≥95% Monomer by analytical SEC, >95% by SDS Page
- Formulation:** This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

Preparation & Storage

- Storage:** Functional grade preclinical antibodies may be stored sterile as received at 2-8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at ≤ -70°C. Avoid Repeated Freeze Thaw Cycles.
Store Bradford Reagent at Room Temperature for 1 Year.
- Shipping:** Next Day 2-8°C
- Preparation:** Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.
- Recommended Dilution Buffer:** In vivo Antibody Diluent pH 7.2

Recommended Usage:	Application	Recommended Usage
	FC	The suggested concentration for this 3G8 antibody for staining cells in flow cytometry is ≤ 2.0 µg per 10 ⁶ cells in a volume of 100 µl or 100µl of whole blood. Titration of the reagent is recommended for optimal performance for each application.

- Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

