

Anti-Mouse CD11c In Vivo Antibody - Low Endotoxin

IVMB0141

Description

This Anti-Mouse CD11c In Vivo Antibody - Low Endotoxin is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

Product Information

SKU:	IVMB0141
Contents:	1mg, 5mg, 25mg, 50mg, 100mg Bradford Reagent: 1 vial (2ml)
Synonyms:	α X Integrin, Integrin α X Chain, CR4, p150, ITGAX
Category:	Monoclonal Antibody
Target:	CD11c
Clone:	N418
Isotype:	IgG
Applications:	CyTOF® FC IHC FF IP PhenoCycler® WB
Specificity:	Clone N418 recognizes an epitope on mouse CD11C.

Antibody Data

Reactivity:	Mouse
Host species:	Armenian Hamster
Expression Host:	-
Immunogen:	Mouse spleen dendritic cells

Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

- Product concentration:** ≥ 5.0 mg/ml
- Endotoxin Level:** < 1.0 EU/mg as determined by the LAL method
- Purity:** $\geq 95\%$ Monomer by analytical SEC, $> 95\%$ by SDS Page
- Formulation:** This monoclonal antibody is aseptically packaged and formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.2 - 7.4 with no carrier protein, potassium, calcium or preservatives added. Due to inherent biochemical properties of antibodies, certain products may be prone to precipitation over time. Precipitation may be removed by aseptic centrifugation and/or filtration.

Preparation & Storage

Storage: Functional grade preclinical antibodies may be stored sterile as received at 2° to 8°C for up to one month. For longer term storage, aseptically aliquot in working volumes without diluting and store at $\leq -70^\circ\text{C}$. Avoid Repeated Freeze Thaw Cycles.
Store Bradford Reagent at Room Temperature for 1 Year.

Shipping: Next Day 2-8°C

Preparation: Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

Recommended Dilution Buffer: In vivo Antibody Diluent pH 7.2

Recommended Usage:	Application	Recommended Usage
	FC	The suggested concentration for this N418 antibody for staining cells in flow cytometry is ≤ 1.0 μg per 10^6 cells in a volume of 100 μl . Titration of the reagent is recommended for optimal performance for each application.
	WB	The suggested concentration for this N418 antibody for use in western blotting is 1-10 $\mu\text{g}/\text{ml}$.

Protein Quantification (Optional): To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol

