

# Mouse Anti-CD11c Low Endotoxin In Vivo Antibody (Phenocycler-Fusion Validated)

IVMB0285

## Description

---

This Mouse Anti-CD11c Low Endotoxin In Vivo Antibody (Phenocycler-Fusion Validated) is supplied as a kit for advanced applications. The kit includes Bradford Reagent to quantify total protein concentration for accurate sample normalization (Optional).

## Product Information

---

<b>SKU:</b>	IVMB0285
<b>Contents:</b>	50ug Bradford Reagent: 1 vial (2ml)
<b>Synonyms:</b>	$\alpha$ X Integrin, Integrin $\alpha$ X Chain, CR4, p150, ITGAX
<b>Category:</b>	Monoclonal Antibody
<b>Target:</b>	CD11c
<b>Clone:</b>	N418
<b>Isotype:</b>	IgG
<b>Applications:</b>	IHC FF PhenoCycler®
<b>Specificity:</b>	Clone N418 recognizes an epitope on mouse CD11C.

## Antibody Data

---

<b>Reactivity:</b>	Mouse
<b>Host species:</b>	Armenian Hamster
<b>Expression Host:</b>	-
<b>Immunogen:</b>	Mouse spleen dendritic cells

### Manufacturers Statement

This final kit system is assembled and quality-released by Assay Genie Limited.

**Product concentration:** 0.5 mg/ml

**Endotoxin Level:** -

**Purity:** -

**Formulation:** This purified antibody is formulated in 0.01 M phosphate buffered saline (150 mM NaCl) PBS pH 7.4.

## Preparation & Storage

---

**Storage:** This antibody is stable for at least one week when stored at 2-8°C. For long term storage, aliquot in working volumes without diluting and store at -20°C in a manual defrost freezer. Avoid Repeated Freeze Thaw Cycles. Store Bradford Reagent at Room Temperature for 1 Year.

**Shipping:** -

**Preparation:** Functional grade preclinical antibodies are manufactured in an animal free facility using in vitro cell culture techniques and are purified by a multi-step process including the use of protein A or G to assure extremely low levels of endotoxins, leachable protein A or aggregates.

**Recommended Dilution Buffer:** -

<b>Recommended Usage:</b>	<b>Application</b>	<b>Recommended Usage</b>
	CODEX®	This CD11c (Clone N418) antibody is formulated to simplify the antibody preparation needed when performing a CODEX® barcode conjugate. The suggested concentration is 0.5 mg/ml.

**Protein Quantification (Optional):** To quantify total protein levels, use the Bradford Reagent included in this kit. Visit <https://www.assaygenie.com/bradford-protein-assay-protocol/> to view the full protocol